

Recombinant SARS-COV-2 Spike RBD(N501Y,K417N,E484K) Protein

Catalog No.: RP02321 Recombinant

Sequence Information

Species Gene ID Swiss Prot SARS-COV-2 QHD43416.1

Tags

C-His

Synonyms

S protein RBD;Spike glycoprotein Receptor-binding domain;S glycoprotein RBD;Spike protein RBD

Product Information

Source

Purification

HEK293 cells

> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC

Endotoxin

< 1 EU/ μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
<u>~</u>	www.abclonal.com.cn

Background

Basic Information

Description

Recombinant SARS-COV-2 Spike RBD (N501Y, K417N,E484K) Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Arg319-Phe541(N501Y,K417N,E484K)) of SARS-COV-2 Spike RBD (N501Y, K417N,E484K) fused with His tag at the C-terminal.

Bio-Activity

Immobilized SARS-CoV-2 Spike RBD (N501Y,K417N,E484K) at 0.5 μ g/mL (100 μ L/Well) on the plate. Dose response curve for Human ACE2, hFc Tag with the EC₅₀ of 0.48 μ g/mL determined by ELISA.

Storage

Store the lyophilized protein at -20°C to -80°C for long term.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



SARS-COV-2 Spike RBD (N501Y,K417N,E484K) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of SARS-COV-2 Spike RBD (N501Y,K417N,E484K) is greater than 95% as determined by SEC-HPLC.



Immobilized SARS-COV-2 Spike RBD (N501Y,K417N,E484K) , His Tag at $0.5\mu g/ml$ (100ul/Well) on the plate. Dose response curve for Human ACE2, hFc Tag with the EC $_{50}$ of 9.9ng/ml determined by ELISA.