

Biotinylated Recombinant SARS-COV-2 Spike S1 Protein

Catalog No.: RP02328 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
SARS-COV-2		QHD43416.1

Tags

C-His-Avi

Synonyms

S1 protein;Spike glycoprotein Subunit1;S glycoprotein Subunit1;Spike protein S1

Product Information

Source	Purification
Expi293	> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening.
Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water.
Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Background

Basic Information

Description

Biotinylated Recombinant SARS-COV-2 SARS-COV-2 Spike S1 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Gln14-Arg683) of SARS-COV-2 SARS-COV-2 Spike S1 fused with His tag and Avi tag at the C-terminal.

Bio-Activity

Immobilized Biotinylated SARS-COV-2 Spike S1,His Tag at 0.5μg/ml (100μl/Well). Dose response curve for Human ACE2,hFc Tag with the EC₅₀ of 16.1ng/ml determined by ELISA.

Storage

Store the lyophilized protein at -20°C to -80°C for long term.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.


Contact

☎ | 400-999-6126


✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn


Validation Data



Biotinylated SARS-COV-2 Spike S1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated SARS-COV-2 Spike S1 is greater than 95% as determined by SEC-HPLC.



Immobilized Biotinylated SARS-COV-2 Spike S1, Hisi Tag at 1μg/ml (100μl/well) on the streptavidin precoated plate (5μg/ml). Dose response curve for Human ACE2, hFc Tag with the EC₅₀ of 13.2ng/ml determined by ELISA.