Biotinylated Recombinant Human IL-10 Protein

Catalog No.: RP02391B Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Human	3586	P22301

Tags

N-His&Avi

Synonyms

CSIF; GVHDS; IL-10; IL10A; TGIF;IL10;GVHDS;IL-10;IL10A;TGIF

Product Information

Source	Purification
HEK293 cells	> 95% by Tris-Bis PAGE

Calculated MW	Observed MW	
21.3 kDa	23-25 kDa	

Endotoxin

< 1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Background

Interleukin 10 (IL10) is a key anti-inflammatory cytokine that can inhibit proinflammatory responses of both innate and adaptive immune cells. An association between IL10 and intestinal mucosal homeostasis became clear with the discovery that IL10 and IL10 receptor (IL10R)-deficient mice develop spontaneous intestinal inflammation. Similarly, patients with deleterious mutations in IL10, IL10RA, or IL10RB present with severe enterocolitis within the first months of life.

Basic Information

Description

Biotinylated Recombinant Human IL-10 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Ser19-Asn178) of Human IL-10 fused with a His tag and Avi tag at the N-terminal.

Bio-Activity

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20 $^{\circ}\text{C}$ for 3 months, at 2-8 $^{\circ}\text{C}$ for up to 1 week.

Avoid repeated freeze/thaw cycles.





Biotinylated Human IL-10 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%. The purity of Biotinylated Human IL-10 is greater than 95% as determined by SEC-HPLC.

Human IL-10 R alpha, His Tag immobilized on CM5 Chip can bind Biotinylated Human IL-10, His Tag with an affinity constant of 0.30 nM as determined in SPR assay (Biacore T200).