

# Biotinylated Recombinant Human IL-17F Protein

Catalog No.: RP02417B **Recombinant**

## Sequence Information

| Species | Gene ID | Swiss Prot |
|---------|---------|------------|
| Human   | 112744  | Q96PD4     |

### Tags

C-His&amp;Avi

### Synonyms

IL17F;CANDF6;IL-17F;ML-1;ML1

## Product Information

| Source       | Purification                             |
|--------------|--|
| HEK293 cells | > 95% by Tris-Bis PAGE;> 95% by SEC-HPLC |

### Endotoxin

&lt; 1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

### Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

The Interleukin 17 (IL-17) family proteins, comprising six members (IL-17A through IL-17F), are secreted, structurally related proteins that share a conserved cystine-knot fold near the C-terminus, but have considerable sequence divergence at the N-terminus. IL-17F is ligand for IL17RA and IL17RC. The heterodimer formed by IL17A and IL17F is a ligand for the heterodimeric complex formed by IL17RA and IL17RC. Involved in stimulating the production of other cytokines such as IL6, IL8 and CSF2, and in regulation of cartilage matrix turnover.

## Basic Information

### Description

Biotinylated Recombinant Human IL-17F Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Arg31-Gln163) of Human IL-17F fused with a His tag and Avi tag at the C-terminal.

### Bio-Activity

### Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

## Validation Data

---



Biotinylated Human IL-17F on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.