

Biotinylated Recombinant Human Low-density lipoprotein receptor/LDLR Protein

Catalog No.: RP02430B **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	3949	P01130

Tags

C-His&Avi

Synonyms

LDLR; FH; FHC; LDLCQ2; low-density lipoprotein receptor;FH;FHC;LDLCQ2

Product Information

Source	Purification
HEK293 cells	> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC

Calculated MW	Observed MW
87.6 kDa	120-140 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

The low density lipoprotein receptor (LDLR) is the founding member of the LDL R family of widely expressed cell surface scavenger receptors. It is a cell-surface receptor that recognizes the apoprotein B100 which is embedded in the phospholipid outer layer of LDL particles.

Basic Information

Description

Biotinylated Recombinant Human Low-density lipoprotein receptor/LDLR Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Ala22-Arg788) of Human LDLR fused with a His tag and Avi tag at the C-terminal.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Biotinylated Human LDLR on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human LDLR is greater than 95% as determined by SEC-HPLC.