

Biotinylated Recombinant Human NKG2-2A/KLRC1/CD159a Protein

Catalog No.: RP02454B **Recombinant**

Sequence Information

Species Human **Gene ID** 3821 **Swiss Prot** P26715

Tags

N-His&Avi

Synonyms

KLRC1;CD159A;NKG2;NKG2A

Background

NKG2, also known as NKG2A(CD159A), is a member of the killer cell lectin-like receptor family. NKG2A marks a unique immune effector subset preferentially co-expressing the tissue-resident CD103 molecule, but not immune checkpoint inhibitors. It is expressed only in NK-cells, but not in T-cells or B-cells.

Basic Information

Description

Biotinylated Recombinant Human NKG2-2A/KLRC1/CD159a Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Arg100-Leu233) of Human NKG2-2A fused with a His tag and Avi tag at the N-terminal.

Bio-Activity

Shipping

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Operational Notes

For your safety and health, please wear a lab coat and disposable gloves for handling.

Product Information

Source

HEK293 cells

Purification

≥ 95 % as determined by Tris-Bis PAGE.

Calculated MW

18.3 kDa

Observed MW

40-50 kDa

Endotoxin

< 1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.


Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.


Contact

	400-999-6126
	cn.market@abclonal.com.cn
	www.abclonal.com.cn

Validation Data



Immobilized Biotinylated Human NKG2A / CD159a,His Tag at 1 μ g/ml (100 μ l/well) on streptavidin precoated (5 μ g/well) plate, can bind NKG2A Ab., with the EC₅₀ of 0.14 μ g/ml determined by ELISA.



Biotinylated Recombinant Human NKG2-2A/KLRC1/CD159a Protein was determined by Tris-Bis PAGE under reducing conditions.