

Biotinylated Recombinant Human ROR2 Protein

Catalog No.: RP02480B Recombinant

Sequence Information

 Species
 Gene ID
 Swiss Prot

 Human
 4920
 A1L4F5(Q019

 74)
 74)

Tags C-His&Avi

Synonyms BDB: BDB1:

NTRKR2;ROR2;BDB1;NTRKR2

Product Information

Source Purification HEK293 cells ≥ 95 % as

K293 cells ≥ 95 % as determined by Tris-Bis PAGE;≥ 95 % as determined by

HPLC.

Calculated MW Observed MW

44.2 kDa 54-65 kDa

Endotoxin

< 1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

ROR2 (Receptor Tyrosine Kinase-like Orphan Receptor 2) is a member of the ROR family of receptor tyrosine kinases and is important for skeletal development, including bone and cartilage formation, as well as for the development of the central nervous system. Mature human ROR2 contains a 369 amino acid (aa) extracellular domain (ECD) and a 518 aa cytoplasmic tail containing an tyrosine kinase domain.

Basic Information

Description

Biotinylated Recombinant Human ROR2 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Val34-Gly403) of Human ROR2 fused with a His and Avi tag at the C-terminal.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80°C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Biotinylated Recombinant Human ROR2 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Biotinylated Human ROR2 is greater than 95% as determined by SEC-HPLC.



Immobilized Anti-ROR2 Antibody, hFc Tag at $0.5\mu g/ml$ (100 $\mu l/well$) on the plate. Dose response curve for Biotinylated Human ROR2, hFc Tag with the EC₅₀ of 6.5ng/ml determined by ELISA.