

# Recombinant Human Siglec-6/CD327 Protein

**Catalog No.: RP02493** Recombinant

## Sequence Information

Species	Gene ID	Swiss Prot
Human	946	O43699-3

### Tags

C-hFc&amp;6\*His

### Synonyms

SIGLEC6; CD33L; CD33L1; OBBP1; Sialic acid-binding Ig-like lectin 6; Siglec-6; CD33 antigen-like 1; CDw327; Obesity-binding protein 1; OB-BP1; CD327

## Product Information

Source	Purification
HEK293 cells	> 95% by Tris-Bis PAGE;> 95% by SEC-HPLC

Calculated MW	Observed MW
58.31 kDa	75-100 kDa

### Endotoxin

&lt;0.1EU/ug of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Background

Siglecs (sialic acid binding Ig-like lectins) are I-type (Ig-type) lectins that belong to the Ig superfamily. They are characterized by an N-terminal Ig-like V-type domain which mediates sialic acid binding, followed by varying numbers of Ig-like C2-type domains. Eleven human Siglecs (Siglec-1 through 11) have been cloned and characterized. Within these eleven, there are at least two groups, one of which is termed the CD33-related group. CD33-related Siglecs include CD33/Siglec-3 and Siglec-5 through 11. To date, no Siglec has been shown to recognize any cell surface ligand other than sialic acid. This suggests that interactions with glycans containing this carbohydrate are important in mediating the biological functions of Siglecs. The cDNA of human Siglec-6 (also known as OB-BP1 and CD33L), encodes a putative 442 amino acid (aa) protein that contains a 15 aa signal peptide, a 321 aa extracellular region, a 21 aa transmembrane region (TM), and an 85 aa cytoplasmic tail. The extracellular region contains one N-terminal V-type Ig-like domain followed by two Ig-like C2-type domains. The cytoplasmic domain has one immunoreceptor tyrosine-based inhibition motif (ITIM). At least three additional isoforms exist, all of which encode an additional 11 aa's at the N-terminus, likely due to the utilization of an alternate start site. Two of the three isoforms also show splicing. One isoform shows a 16 aa in-frame deletion in the second C2-like domain, while the other shows a deletion of the TM and cytoplasmic region, thus potentially generating a soluble form.

## Basic Information

### Description

Recombinant Human Siglec-6/CD327 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln27-Val331) of Human Siglec-6/CD327 (Accession #NP\_942142.3) fused with a hFc and His tag at the C-terminus.

### Bio-Activity

### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## Contact

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)
 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Validation Data

---



Recombinant Human Siglec-6/CD327 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 75-100kDa.