

Recombinant Human Siglec-10 Protein

Catalog No.: RP02498 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	89790	Q96LC7

Tags

C-hFc

Synonyms

SLG2; PRO940; SIGLEC-10; SIGLEC10

Product Information

Source	Purification
HEK293 cells	> 95% by Tris-Bis PAGE

Endotoxin

< 0.1 EU/μg

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact

 | 400-999-6126 | cn.market@abclonal.com.cn | www.abclonal.com.cn

Background

Siglec-10 is a ligand for CD52, the target of the therapeutic monoclonal antibody Alemtuzumab. It is also reported to bind to Vascular adhesion protein 1 (VAP-1) and to the co-stimulatory molecule CD24 also known as HSA (Heat-stable antigen). Siglecs (sialic acid binding Ig-like lectins) are I-type lectins that belong to the immunoglobulin superfamily. They are characterized by an N-terminal Ig-like V-type domain which mediates sialic acid binding, followed by a varying number of Ig-like C2-type domains. Siglecs 5-11 constitute the CD33/Siglec-3 related group, and are differentially expressed in the hematopoietic system.

Basic Information

Description

Recombinant Human Siglec 10 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Met17-Thr546) of Human Siglec 10 fused with hFc tag at the C-terminal.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human Siglec-10 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 100-140 kDa.