

Recombinant Human IL-37/IL-1F7 Protein

Catalog No.: RP02509 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 27178 O9NZH6

Tags

C-His

Synonyms

IL37;FIL1;FIL1(ZETA);FIL1Z;IL-1F7;IL-1H;I L-1H4;IL-1RP1;IL-37;IL1F7;IL1H4;IL1RP1

Product Information

Source Purification *E. coli* > 95% by SDS-PAGE.

Endotoxin

<0.1EU/µg

Formulation

Lyophilized from a 0.2 μ m filtered solution of PBS, pH 8.0.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

Decreasing synthesis of IL-37 in PBMCs will cause increasing production of proinflammatory cytokines such as IL-1 alpha, IL-1 beta, IL-6 and TNF- alpha. The IL-37 work as an innate inflammatory inhibitor is also proved by its high expression in synovial tissue from rheumatoid arthritis patients. Full length IL-37 locates mainly in the cytoplasm, and it can translocate to the nucleus after activation through cleavage by CASP1, then exerts its activity by directly interacting with SMAD3.

Basic Information

Description

Recombinant Human IL-37 Protein is produced by Escherichia coli expression system. The target protein is expressed with sequence of Human IL-37 (Accession #NP 775297.1) fused with His at C terminus.

Bio-Activity


Measured by its ability to inhibit IL-8 secretion in human PBMCs. The ED $_{50}$ for this effect is <0.9 $\mu g/mL$.

Storage

Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human IL-37/IL-1F7 Protein was determined by SDS-PAGE with Coomassie Blue, showing bands at 20 kDa.