

Catalog No.: RP02521 **Recombinant**

Species	Gene ID	Swiss Prot
Mouse	16173	P70380

C-His

Synonyms
lg; ll-; lgif; ll-18; LL18

Source	Purification
HEK293 cells	≥ 95 % as determined by SDS-PAGE

18.96 kDa 15-25 kDa

< 0.1 EU/μg of the protein by LAL method.

Lyophilized from a 0.22 μm filtered solution of PBS, pH 8.0.

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Interleukin-18 (IL-18) is a cytokine that belongs to the IL-1 superfamily and is produced by macrophages and other cells. IL-18 works by binding to the interleukin-18 receptor, and together with IL-12 it induces cell-mediated immunity following infection with microbial products like lipopolysaccharide (LPS). After stimulation with IL-18, natural killer (NK) cells and certain T cells release another important cytokine called interferon- γ (IFN- γ) or type II interferon that plays an important role in activating the macrophages or other cells. The combination of this cytokine and IL-12 has been shown to inhibit IL-4 dependent IgE and IgG1 production, and enhance IgG2a production in B cells. IL-18 binding protein (IL-18BP) can specifically interact with this cytokine, and thus negatively regulate its biological activity.

Recombinant Mouse IL-18 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence of Mouse IL-18 fused with His tag at the C-terminus

Measured by its ability to induce IFN-gamma secretion by KG1 human acute myelogenous leukemia cells in the presence of TNF-alpha (Catalog: RP00001). The ED₅₀ for this effect is 42.93-171.72 ng/mL, corresponding to a specific activity of 0.58×10⁴~2.3×10⁵ units/mg.


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.

 | 400-999-6126


 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Validation Data



Recombinant Mouse IL-18 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Mouse IL-18 induce IFN-gamma secretion by KG1 human acute myelogenous leukemia cells in the presence of TNF-alpha(Catalog: RP00001). The ED₅₀ for this effect is 42.93-171.72 ng/mL, corresponding to a specific activity of 0.58×10⁴~2.3×10⁵ units/mg.