

Recombinant Human FGFR-2 alpha (IIIb)/KGFR/CD332 Protein

Catalog No.: RP02530 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 2263 P21802-3

Tags C-hFc

Synonyms

FGF R2a; FGFR2 alpha; KGFR; CD332; BBDS; BEK; BFR-1; CEK3; CFD1; ECT1; JWS; K-SAM; KGFR; TK14; TK25; FLJ98662

Product Information

Source Purification HEK293 cells ≥ 95 % as

determined by Tris-Bis PAGE;≥ 95 % as determined by

HPLC.

Calculated MW Observed MW

66.4 kDa 80-115 kDa

Endotoxin

< 1 EU/ μg of the protein by LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>		400-999-6126
\bowtie		cn.market@abclonal.com.cn
•	Τ	www.abclonal.com.cn

Background

Four distinct genes encoding closely related FGF receptors, FGF R1 - 4, are known. All four genes for FGF Rs encode proteins with an N-terminal signal peptide, three immunoglobulin (Ig)-like domains, an acid-box region containing a run of acidic residues between the IgI and IgII domains, a transmembrane domain and the split tyrosine-kinase domain. Multiple forms of FGF R1 - 3 are generated by alternative splicing of the mRNAs. A frequent splicing event involving FGF R1 and 2 results in receptors containing all three Ig domains, referred to as the alpha? isoform, or only IgII and IgIII, referred to as the beta? isoform.

Basic Information

Description

Recombinant Human FGFR-2 alpha (IIIb)/KGFR/CD332 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Arg22-Glu378) of human FGFR-2 alpha (IIIb)/KGFR/CD332 (Accession #P21802-3) fused with hFc tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Recombinant Human FGFR-2 alpha (IIIb)/KGFR/CD332 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Human FGFR2 alpha (IIIb) is greater than 95% as determined by SEC-HPLC.



Immobilized Human FGFR2 alpha IIIb, hFc tag at $0.5\mu g/ml$ (100 $\mu l/Well$). Dose response curve for Biotinylated Anti-FGFR2 Ab., hFc Tag with the EC $_{50}$ of 10.0ng/ml determined by ELISA.