

# Recombinant Human FGFR2 alpha (IIIb)/KGFR/CD332 Protein

Catalog No.: RP02533 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	2263	P21802-3

### Tags

C-His

### Synonyms

FGF R2a; FGFR2 alpha; KGFR; CD332; BBDS; BEK; BFR-1; CEK3; CFD1; ECT1; JWS; K-SAM; KGFR; TK14; TK25; FLJ98662

## Product Information

### Source

HEK293 cells

### Purification

> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

### Endotoxin

Less than 1EU per µg by the LAL method.

### Formulation

### Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

☎ | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

🌐 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

Four distinct genes encoding closely related FGF receptors, FGF R1 - 4, are known. All four genes for FGF Rs encode proteins with an N-terminal signal peptide, three immunoglobulin (Ig)-like domains, an acid-box region containing a run of acidic residues between the IgI and IgII domains, a transmembrane domain and the split tyrosine-kinase domain. Multiple forms of FGF R1 - 3 are generated by alternative splicing of the mRNAs. A frequent splicing event involving FGF R1 and 2 results in receptors containing all three Ig domains, referred to as the alpha? isoform, or only IgII and IgIII, referred to as the beta? isoform.

## Basic Information

### Description


Recombinant Human FGFR2 alpha (IIIb) Protein is expressed from Expi293 with His tag at the C-terminal. It contains Arg22-Glu378.

### Bio-Activity


### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


## Validation Data




Human FGFR2 alpha (IIIb) on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.



The purity of Human FGFR2 alpha (IIIb) is greater than 95% as determined by SEC-HPLC.



Immobilized Human FGFR2 alpha (IIIb) , His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-FGFR2 alpha (IIIb) Antibody, hFc Tag with the  $EC_{50}$  of 10.4ng/ml determined by ELISA.



Serial dilutions of Anti-FGFR2 alpha (IIIb) Antibody were added into Human FGFR2 alpha (IIIb) , His Tag : Biotinylated Human FGF10, No Tag binding reactions.