

Biotinylated Recombinant Human BTN1A1/Butyrophilin Protein

Catalog No.: RP02557 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	696	Q13410

Tags

C-His&Avi

Synonyms

BT; BTN1A1; BTN; Butyrophilin

Product Information

Source

HEK293 cells

Purification

> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Endotoxin


Less than 1EU per µg by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

Butyrophilin 1A1 (BTN1A1) is one of the membrane proteins that surrounds LD. BTN1A1 plays an important role in regulating LD synthesis via a mechanism involving membrane phospholipid composition.

Basic Information

Description

Recombinant Biotinylated Human BTN1A1/Butyrophilin Protein is expressed from Expi293 with His tag and Avi tag at the C-terminal. It contains Ala27-Arg242.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for long term.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Biotinylated Human BTN1A1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Biotinylated Human BTN1A1 is greater than 95% as determined by SEC-HPLC.