

# Recombinant Cynomolgus 5'-Nucleotidase/NT5E/CD73 Protein

Catalog No.: RP02630 Recombinant

## **Sequence Information**

**Species Gene ID Swiss Prot** Cynomolgus EHH53214.1

**Tags** C-His

**Synonyms** 

CD73; NT5E; NTE; 5' nucleotidase (CD73); 5'-NT; 5-NT; E5NT; EC 3.1.3.5; eN; eNT; NT; NT55'-nucleotidase; NT5; NTE; CALJA

## **Product Information**

Source Purification

HEK293 cells > 95% as

determined by Tris
Ris PAGE□> 95% as

Bis PAGE□> 95% as determined by HPLC

Calculated MW Observed MW

59.6 kDa 60-70 kDa

**Endotoxin** 

Less than 1EU per  $\mu g$  by the LAL method.

#### **Formulation**

## Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
<u>~</u>	T	www.abclonal.com.cn

## **Background**

CD73, also known as ecto-5'-nucleotidase, is an enzyme that in humans is encoded by the NT5E gene.CD73 commonly serves to convert AMP to adenosine.The enzyme consists of a dimer of 2 identical 70-kD subunits bound by a glycosyl phosphatidyl inositol linkage to the external face of the plasma membrane. The enzyme is used as a marker of lymphocyte differentiation. A? deficiency of CD73 occurs in a variety of immunodeficiency diseases.

## **Basic Information**

## **Description**

Recombinant Cynomolgu 5'-Nucleotidase/NT5E/CD73 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Trp27-Cys554) of Cynomolgus 5'-Nucleotidase/NT5E/CD73 fused with His tag at the C-terminus.

## **Bio-Activity**


#### Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20  $^{\circ}$ C for 3 months, at 2-8  $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.


## **Validation Data**



Cynomolgus CD73 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Cynomolgus CD73 is greater than 95% as determined by SEC-HPLC.



Immobilized Cynomolgus CD73, His Tag at  $1\mu g/ml$  (100 $\mu l/Well)$  on the plate. Dose response curve for Anti-CD73 Antibody, hFc Tag with the EC $_{50}$  of 10.8ng/ml determined by ELISA.