

Recombinant Cynomolgus B7-2/CD86 Protein

Catalog No.: RP02634 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Cynomolgus		G7NXR4

Tags

C-His

Synonyms

CD86 molecule; CD86; B70; B7-2 antigen; B72; B7-2; BU63; FUN-1; LAB72; MGC34413; CD28LG2

Product Information

Source	Purification
HEK293 cells	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Endotoxin

Less than 1EU per µg by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize freeze-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

B7-1 and B7-2 are homologous costimulatory ligands expressed on the surface of antigen presenting cells (APCs). Binding of these molecules to the T cell costimulatory receptors, CD28 and CTLA-4, is essential for the activation and regulation of T cell immunity. B7-1 and B7-2 do not form hetero-oligomers, underscoring the biological relevance of dimeric and monomeric state of B7-1 and B7-2, respectively.

Basic Information

Description


Recombinant Cynomolgus B7-2/CD86 Protein is expressed from Expi293 with His tag at the C-terminal. It contains Tyr26-His240.

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for long term. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data



Cynomolgus B7-2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Cynomolgus B7-2 is greater than 95% as determined by SEC-HPLC.



Cynomolgus B7-2, His Tag immobilized on CM5 Chip can bind Cynomolgus CTLA-4, His Tag with an affinity constant of $0.102 \mu\text{M}$ as determined in SPR assay (Biacore T200).