Recombinant Cynomolgus B7-H2/ICOSLG/CD275 Protein

Catalog No.: RP02644 Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Cynomolgus	102132485	XP_00554861
		7.1

Tags C-His

Synonyms

CD275; B7H2; B7-H2; B7RP-1; GL50; ICOSL; ICOS-L; ICOSLG; B7RP1; LICOS; ICOS ligand; B7-H2B7-related protein 1; B7RP-1; B7RP1B7-like protein GI50

Product Information

Source Purification HEK293 cells ≥ 95 % as determined by Tris-Bis PAGE;≥ 95 % as

determined by HPLC.

50-70 kDa

Calculated MW Observed MW

27.7 kDa

Endotoxin

< 1 EU/ μ g of the protein by LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Background

B7-H2, also known as B7-related protein (B7RP1), ICOS Ligand, and CD275, is an approximately 60 kDa transmembrane glycoprotein in the B7 family of immune regulatory molecules. B7-H2 is a ligand for the T-cell-specific cell surface receptor ICOS. Acts as a costimulatory signal for T-cell proliferation and cytokine secretion; induces also B-cell proliferation and differentiation into plasma cells.

Basic Information

Description

Recombinant CynomolgusB7-H2/ICOSLG/CD275 Protein is produced by Expi293 expression system. The target protein is expressed with sequence (Asp19-Thr256) of Cynomolgus B7-H2 fused with His tag at the C-terminus.

Bio-Activity


Storage


Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.







The purity of Cynomolgus B7-H2 is greater than 95% as determined by SEC-HPLC.

Human ICOS, hFc Tag captured on CM5 Chip via Protein A can bind Cynomolgus B7-H2, His Tag with an affinity constant of 5.42 nM as determined in SPR assay (Biacore T200).

Recombinant Cynomolgus B7-

H2/ICOSLG/CD275 Protein was determined by Tris-Bis PAGE under reducing conditions.