

Recombinant Cynomolgus Guanylyl cyclase C/GUCY2C Protein

Catalog No.: RP02645 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Cynomolgus	102130850	XP_00557027 0.1

Tags

C-His

Synonyms

Guanylyl cyclase C; GC-C; STAR;
GUCY2C; GUC2C; STA receptor; DIAR6;
EC 4.6.1; GCC; GUC2CEC 4.6.1.2; MUCIL

Product Information

Source	Purification
HEK293 cells	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Endotoxin

Less than 1EU per µg by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening.
Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

		400-999-6126
		cn.market@abclonal.com.cn
		www.abclonal.com.cn

Background

Guanylyl cyclase C (GUCY2C) has canonical centrality in defense of key intestinal homeostatic mechanisms, encompassing fluid and electrolyte balance, epithelial dynamics, antitumorigenesis, and intestinal barrier function. GUCY2C may represent a new target for anti-obesity pharmacotherapy.

Basic Information

Description


Recombinant Cynomolgus Guanylyl cyclase C/GUCY2C Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser24-Gln430) of cynomolgus Guanylyl cyclase C/GUCY2C (Accession #XP_005570270.1) fused with His tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.


Validation Data



Cynomolgus GUCY2C on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Cynomolgus GUCY2C is greater than 95% as determined by SEC-HPLC.



Immobilized Cynomolgus GUCY2C, His Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Anti-GUCY2C Antibody, hFc Tag with the EC_{50} of 103.3ng/ml determined by ELISA.