

Recombinant Human EPHB2 Protein

Catalog No.: RP02657 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	2048	P29323-1

Tags

C-His

Synonyms

DRT; ERK; HEK5; TYRO5; CAPB; EK5; EPHT3; PCBC; Cek5; EphB2; EPHT3MGC87492; Nuk; Qek2; Sek3

Product Information

Source	Purification
HEK293 cells	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Calculated MW	Observed MW
59.2 kDa	65-75 kDa

Endotoxin

Less than 1EU per µg by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

EphB2, a receptor tyrosine kinase for ephrin ligands, is overexpressed in various cancers and plays an important role in tumor progression. EPHB2 promotes endothelial-mesenchymal transition (EMT) and elicits associated pathologic characteristics of glioblastoma multiforme (GBM) such as invasion and migration. EPHB2 is epigenetically overexpressed in hypoxia, a condition highly prevalent in malignancy. Furthermore, HIF-2α is required for EPHB2 stabilization by hypoxia.

Basic Information

Description


Recombinant Human EPHB2 Protein is expressed from Expi293 with His tag at the C-terminal. It contains Val19-Leu543.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data



Human EPHB2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Human EPHB2 is greater than 95% as determined by SEC-HPLC.



Immobilized Human EPHB2, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-EPHB2 Antibody, hFc Tag with the EC₅₀ of 14.9ng/ml determined by ELISA.