

Recombinant Cynomolgus GARP&Latent TGF beta 1 Complex Protein

Catalog No.: RP02690 Recombinant

Sequence Information

Species Gene ID Cynomolgus

Swiss Prot A0A2K5X2X9

(GARP)&A0A 2K5TJB2(Late nt TGF beta

Tags

C-His&Avi

Synonyms

LRRC32; GARP; LAP; TGF-beta-1; LRRC32&TGF-beta 1; LRRC32&TGFB1

Product Information

Source

Purification

HEK293 cells > 95% as determined

determined by Tris-Bis PAGE□> 95% as determined by HPLC

Endotoxin

Less than 1EU per µg by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid votex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

GARP&Latent TGF Beta is a complex found on surface of many types of cells. In Tregs, GARP is involved in TCR-mediated activation of Latent TGF- β and thus promoting secretion and activation of TGF- β . Integrin $\alpha\nu\beta8$ on the surface of immune cells and other cells recognizes RGD in LAP, resulting in the release of mature TGF- β from the TGF- β &GARP complex.

Basic Information

Description

Recombinant Cynomolgus GARP&Latent TGF beta 1 Complex Protein is expressed from Expi293 with His tag and Avi tag at the C-terminal. [It contains Ala18-Leu628(GARP)&Leu30-Ser390(Latent TGF beta 1).

Bio-Activity


Storage

Store the lyophilized protein at -20°C to -80°C for long term.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data



Cynomolgus GARP&Latent TGF beta 1 Complex on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Cynomolgus GARP&Latent TGF beta 1 Complex is greater than 95% as determined by SEC-HPLC.



Immobilized Cynomolgus GARP&Latent TGF beta 1 Complex, His Tag at $2\mu g/ml$ (100 $\mu l/Well$) on the plate. Dose response curve for Anti-GARP&Latent TGF Beta 1 Antibody, hFc Tag with the EC $_{50}$ of 11.0ng/ml determined by ELISA.