

Recombinant Human KIR3DL2/CD158k Protein

Catalog No.: RP02706 **Recombinant**

Sequence Information

| Species | Gene ID | Swiss Prot |
|---------|---------|------------|
| Human | 3812 | P43630-1 |

Tags

C-His&Avi

Synonyms

NKAT-4; NKAT4; CD158k; CL-5; KIR3DL2; NKAT4A; NKAT4B; p140

Product Information

| Source | Purification |
|--------------|--|
| HEK293 cells | > 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC |

Endotoxin

Less than 1EU per µg by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Background

KIR3DL2 is a member of the killer cell immunoglobulin-like receptor (KIR) family that was initially identified at the surface of natural killer (NK) cells. KIR3DL2, also known as CD158k, is expressed as a disulfide-linked homodimer. Each chain is composed of three immunoglobulin-like domains and a long cytoplasmic tail containing two immunoreceptor tyrosine-based inhibitory motifs.

Basic Information

Description


Recombinant Human KIR3DL2/CD158k Protein is produced by Expi293 cells expression system. The target protein is expressed with sequence (Leu22-Leu339) of Human KIR3DL2/CD158k (Accession #P43630-1) fused with His tag and Avi tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Validation Data



Human KIR3DL2 on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.



The purity of Human KIR3DL2 is greater than 95% as determined by SEC-HPLC.



Immobilized Human KIR3DL2, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Anti-KIR3DL2 Antibody, hFc Tag with the EC₅₀ of 0.47µg/ml determined by ELISA.