

Recombinant Mouse NKG2-D/KLRK1/CD314 Protein

Catalog No.: RP02720 Recombinant

Sequence Information

Species Gene ID Swiss ProtMouse 27007 054709-1

Tags

N-hFc

Synonyms

CD314; D12S2489E; KLR; NKG2-D; NKG2D

Product Information

Source

Purification

HEK293 cells

> 95% as determined by Tris-Bis PAGE∏> 95% as determined by HPLC

Calculated MW Observed MW

Endotoxin

<0.1EU/µg by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
$lue{f \odot}$	www.abclonal.com.cn

Background

NKG2D is a type II transmembrane glycoprotein having an extracellular lectin-like domain. This domain lacks the recognizable calcium-binding sites found in true C-type lectins and binds protein rather than carbohydrate ligands. Human NKG2D is expressed on CD8 alpha beta T cells, gamma δ T cells, NK cells and NKT cells.

Basic Information

Description

Recombinant Mouse NKG2-D/KLRK1/CD314 Protein is expressed by HEK293 cells expression system. The target protein is expressed with sequence Phe90-Val232 of NKG2D/CD314 (Accession #NP_149069.1) fused with a hFc at the N-terminus

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Mouse NKG2D/CD314 Protein was resolved with SDS PAGE under reducing (R) and non-reducing (NR) conditions [showing single bands at 50-60 kDa and 100-140 kDa, respectively.



The purity of Mouse NKG2D is greater than 95% as determined by SEC-HPLC.