

Recombinant Cynomolgus TNFSF7/CD27 Ligand/CD70 Protein

Catalog No.: RP02728 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Cynomolgus	102130438	G7PYU6-1

Tags

N-His

Synonyms

CD70 molecule; CD70; TNFSF7; CD27 Ligand; CD27-L; CD27LG;?Ki-24 antigen; TNFSF7G

Product Information

Source	Purification
HEK293 cells	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Calculated MW	Observed MW
18.4 kDa	55-70 kDa

Endotoxin

Less than 1EU per µg by the LAL method.

Formulation

Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

CD70, also named CD27 ligand (CD27L), is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and has been designated TNFSF7. CD70 is a cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.

Basic Information

Description

Recombinant Cynomolgus TNFSF7/CD27 Ligand/CD70 Protein is produced by Expi293 cells expression system. The target protein is expressed with sequence (Gln39-Pro194) of Cynomolgus TNFSF7 (Accession # G7PYU6-1) fused with His tag at the N-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.


After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Cynomolgus CD27 Ligand on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



Immobilized Cynomolgus CD27 Ligand, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Cynomolgus/Rhesus macaque CD27, hFc Tag with the EC₅₀ of 41.5ng/ml determined by ELISA.