Recombinant Human DMBT1 Protein



Catalog No.: RP02790 Recombinant

Sequence Information

Species Gene ID Swiss Prot Human 1755 Q9UGM3-2

Tags C-His

Synonyms GP340; DMBT1

Product Information

Source HEK293 cells Purification ≥ 95 % as

determined by SDS-

PAGE.

Calculated MW Observed MW

22.6 kDa 35-45 kDa

Endotoxin

< 0.1 EU/ μ g of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

<u>a</u>	400-999-6126
\bowtie	cn.market@abclonal.com.cn
•	www.abclonal.com.cn

Background

Deleted in malignant brain tumors 1 protein, also known as glycoprotein 34, surfactant pulmonary-associated D-binding protein, DMBT1 and GP34, is a secreted protein which belongs to the DMBT1 family. DMBT1 contains 2CUB domains, 14SRCR domains and 1ZP domain. It is highly expressed in alveolar and macrophage tissues. In some macrophages, expression is detected on the membrane, and in other macrophages, it is strongly expressed in the phagosome/phagolysosome compartments. Defects in DMBT1 are involved in the development of glioma (GLM). Gliomas are central nervous system neoplasms derived from glial cells and comprise astrocytomas, glioblastoma multiforme, oligodendrogliomas, and ependymomas. DMBT1 may be considered as a candidate tumor suppressor for brain, lung, esophageal, gastric, and colorectal cancers. It may play roles in mucosal defense system, cellular immune defense and epithelial differentiation. DMBT1 may play a role as an opsonin receptor for SFTPD and SPAR in macrophage tissues throughout the body, including epithelial cells lining the gastrointestinal tract. It may be an important factor in fate decision and differentiation of transit-amplifying ductular (oval) cells within the hepatic lineage. DMBT1 may function as a binding protein in saliva for the regulation of taste sensation. It binds to HIV-1 envelope protein and has been shown to both inhibit and facilitate viral transmission.

Basic Information

Description

Recombinant Human DMBT1 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Met1-Ser220) of human DMBT1 (Accession #NP_004397.2) fused with a 6×His tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20 $^{\circ}$ C for 3 months, at 2-8 $^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human DMBT1 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.