Recombinant Mouse CXCL9/MIG Protein

Catalog No.: RP02851 Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
Mouse	17329	P18340
Tawa		

Tags C-His

Synonyms

Mig; Scyb9

Product Information

Source	Purification
Pichia	> 95% by SDS
	PAGE

SDS-PAGE.

Endotoxin

Please contact us for more information.

Formulation

Lyophilized from sterile PBS, pH 7.4

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

6	400-999-6126
\times	cn.market@abclonal.com.cn
€	www.abclonal.com.cn

Background

Chemokine (C-X-C motif) ligand 9 (CXCL9), also known as Monokine induced by gamma interferon (MIG), is a small cytokine belonging to the CXC chemokine family. The function of this chemokine has not been specifically defined; however, it is thought to be involved in T cell trafficking. CXCL9/MIG functions as one of the three ligands of chemokine receptor CXCR3 which is a G protein-coupled receptor found predominantly on T cells. CXCL9/MIG, together with CXCL10 and CXCL11, may activate CXCR3 by binding to it. CXCL9 serves as a cytokine that affects the growth, movement, or activation state of cells that participate in immune and inflammatory response. It has been observed that tumour endothelial cells secrete high levels of CXCL9 in all, and CXCL10 in most melanoma metastases. Experiment data represent novel mechanisms by which tumour cells in melanoma metastases might use the chemokine-expressing endothelium to leave the tumour and eventually to form additional metastases at distinct sites. Experiment results also improved that CXCL9/MIG plays an important role in CD4+ T lymphocyte recruitment and development of CAV, MOMA-2+ macrophages are the predominant recipient-derived source of CXCL9/MIG, and recipient CD4 lymphocytes are necessary for sustained CXCL9/MIG production and CAV development in this model. Neutralization of the chemokine CXCL9/MIG may have the rapeutic potential for the treatment of chronic rejection after heart transplantation.

Basic Information

Description

Recombinant Mouse CXCL9/MIG Protein is produced by Pichia expression system. The target protein is expressed with sequence (Thr 22-Thr 126) of mouse CXCL9/MIG (Accession #NP_032625.2) fused with a His tag at the C-terminus..

Bio-Activity

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.





Recombinant Mouse CXCL9/MIG Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 12-15 kDa.