# Recombinant Human GTPase KRas/KRAS (Q61H) Protein

Catalog No.: RP02979LQ Recombinant

### **Sequence Information**

Background

SpeciesGene IDHuman3845

Swiss Prot P01116-2

**Tags** N-8His

#### Synonyms

NŠ; NS3; OES; CFC2; RALD; K-Ras; KRAS1; KRAS2; RASK2; KI-RAS; C-K-RAS; K-RAS2A; K-RAS2B; K-RAS4A; K-RAS4B; K-Ras 2; 'C-K-RAS; c-Ki-ras; c-Ki-ras2

## **Product Information**

| Source  | Purification       |
|---------|--------------------|
| E. coli | ≥ 95 % as          |
|         | determined by SDS- |
|         | PAGE.              |
|         |                    |

# Calculated MW Observed MW 23.5 kDa 25-28 kDa

#### Endotoxin

Please contact us for more information.

#### Formulation

Supplied as a 0.22  $\mu m$  filtered solution in 20mM Tris, 150mM NaCl, 1mM DTT, 10% glycerol, pH7.4

#### Reconstitution

# Contact

| 6        | 400-999-6126              |
|----------|---------------------------|
| $\times$ | cn.market@abclonal.com.cn |
| €        | www.abclonal.com.cn       |

# **Basic Information**

#### Description

Recombinant Human GTPase KRas/KRAS (Q61H) Protein is produced by *E. coli* expression system. The target protein is expressed with sequence of Human GTPase KRas/KRAS (Q61H) (Accession #)fused with N-terminal 8x his tag+ TEV cleavage site and C-terminal avi tag.

#### **Bio-Activity**


KRAS-Q61H activity test using HTRF method.

#### Storage


Store at -70°C. This product is stable at  $\leq$  -70°C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature. Avoid repeated freeze-thaw cycles. Avoid repeated freeze/thaw cycles.




### **Validation Data**



Recombinant Human GTPase KRas/KRAS (Q61H) Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



KRAS-Q61H activity test using HTRF method. The KRAS-Q61H activity was assayed with HTRF technology. The reaction was performed by incubating the KRAS-Q61H protein, GTP, cRAF and beads at 25°C for 60 min, then reading Ratio 665/620nm signal with BMG.



KRAS-Q61H activity test using HTRF method. The KRAS-Q61H activity was assayed with HTRF technology. The reaction was performed by incubating the KRAS-Q61H protein, GTP, cRAF and beads at 25°C for 60 min, then reading Ratio 665/620nm signal with BMG.