

Recombinant Human ING4 Protein

Catalog No.: RP03219 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	51147	Q9UNL4-1

Tags

N-His

Synonyms

ING4; my036; p29ING4

Product Information

Source	Purification
E.coli	> 85% by SDS-PAGE.

Endotoxin

Please contact us for more information.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

ING4 is similar to ING1, a tumor suppressor protein that can interact with TP53, inhibit cell growth, and induce apoptosis. ING4 contains a PHD-finger, which is a common motif in proteins involved in chromatin remodeling. ING4 protein can bind TP53 and EP3/p3, a component of the histone acetyltransferase complex, suggesting its involvement in the TP53-dependent regulatory pathway. ING4 is a component of the HBO1 complex which has a histone H4-specific acetyltransferase activity, a reduced activity toward histone H3 and is responsible for the bulk of histone H4 acetylation in vivo. Through chromatin acetylation, it may function in DNA replication. ING4 may also inhibit tumor progression by modulating the transcriptional output of signaling pathways that regulate cell proliferation.

Basic Information

Description


Recombinant Human ING4 Protein is produced by E.coli expression system. The target protein is expressed with sequence (Met1-Lys249) of human ING4 (Accession #Q9UNL4-1) fused with His tag at the N-terminus.

Bio-Activity

Storage

Store the lyophilized protein at -20°C to -80 °C for twelve months from date of receipt. After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week. Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human ING4 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 25-30 kDa.