

# Recombinant Human HGF Activator/HGFA Protein

Catalog No.: RP03236 **Recombinant**

## Sequence Information

Species	Gene ID	Swiss Prot
Human	3083	Q04756

### Tags

C-His

### Synonyms

HGF activator; HGFA; HGFAC

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

### Endotoxin

&lt; 1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening.  
Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water.  
Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Contact

 | 400-999-6126 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn) | [www.abclonal.com.cn](http://www.abclonal.com.cn)

## Background

HGF activator (HGFA) is a serum-derived serine protease and belongs to the peptidase family S1. HGFA is responsible for the conversion of hepatocyte growth factor (HGF), from the inactive single-chain precursor to the active heterodimeric form, which is a potent mitogen, motogen, and morphogen for liver cells, epithelial cells, and endothelial cells. HGFA is synthesized and secreted by the liver and circulates in the plasma as an inactive single-chain zymogen in normal states. The zymogen is cleaved by thrombin or thermolysin through the endoproteolytic process and forms an active heterodimer linked by a disulfide bond. In turn, the active protease can be inhibited by HGFA inhibitors (HAIs) including HAI-1 and HAI-2. Besides, the HGFA zymogen acquires a strong affinity upon activation and thus may ensure the local action in tissue regeneration in the liver, kidney, and skin. It has been reported that activation of HGF is a critical limiting step in the HGF/SF-induced signaling pathway mediated by Met, and accordingly, aberrant expression of HGFA is implicated in tumorigenesis and progression.

## Basic Information

### Description

Recombinant Human HGF Activator/HGFA Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln36-Ser655) of Human HGF Activator/HGFA (Accession #NP\_001519.1) fused with His tag at the C-terminus.

### Bio-Activity


### Storage

Store the lyophilized protein at -20°C to -80°C for 12 months.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human HGF Activator/HGFA Protein was determined by SDS-PAGE with Coomassie Blue.