

Recombinant Human CDK1&Cyclin B1 Protein

Catalog No.: RP03302LQ **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
Human	983&891	P06493&P14635

Tags

N-GST (CDK1) & N-Flag-His (Cyclin B1)

Synonyms

CDK1; CDC2; CDC28A; CDKN1; P34CDC2; Cyclin-dependent kinase 1; Cyclin B1; CCNB1; CCNB; G2/mitotic-specific cyclin-B1

Product Information

Source	Purification
Baculovirus-Insect Cells	> 90% by SDS-PAGE and HPLC

Calculated MW	Observed MW
60.6 kDa/33.8 kDa	50-60 kDa/28-33 kDa

Endotoxin

< 1.0 EU/μg of the protein by LAL method

Formulation

Supplied as a 0.22 μm filtered solution in 50 mM Tris-HCl, 200 mM NaCl, 20% glycerol, 1 mM DTT. (pH 7.5). Contact us for customized product form or formulation.

Reconstitution

Please use running water to thaw it quickly.

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Background

CDK1 also known as cyclin-dependent kinase 1 or cell division cycle protein 2 homolog is a highly conserved protein that functions as a serine/threonine protein kinase, and is a key player in cell cycle regulation. CDK1 is a small protein (approximately 34 kDa), and is highly conserved. When bound to its cyclin partners, CDK1 phosphorylation leads to cell cycle progression. Given its essential role in cell cycle progression, CDK1 is highly regulated. Most obviously, CDK1 is regulated by its binding with its cyclin partners. Cyclin binding alters access to the active site of CDK1, allowing for CDK1 activity. Cyclin B1 contributes to the switch-like all or none behavior of the cell in deciding to commit to mitosis. Cyclin B1-CDK1 is involved in the early events of mitosis, such as chromosome condensation, nuclear envelope breakdown, and spindle pole assembly. The role of cyclin B1 is to transition the cell from G2 to M phase but becomes unregulated in cancer cells where overexpression of cyclin B1 can lead to uncontrolled cell growth by binding to its partner Cdk. Binding of Cdk can lead to phosphorylation of other substrates at inappropriate time and unregulated proliferation.

Basic Information

Description

Recombinant Human CDK1&Cyclin B1 Protein is produced by Baculovirus-Insect Cells expression system. The target protein is expressed with sequence (Met1-Met297 (CDK1) & Asn165-Val433 (Cyclin B1)) of Human CDK1&CCNB1 (Accession #P06493&P14635) fused with a N-GST (CDK1) & N-Flag-His (Cyclin B1) tag.

Bio-Activity

The activity of CDK1/Cyclin B1 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.


Storage

Store at -70°C. This product is stable at ≤ -70°C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature.


Aliquots below 10 μL are not advisable. Product must not be stored in diluted solutions. Avoid repeated freeze-thaw cycles.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human CDK1&Cyclin B1 Protein was resolved with SDS-PAGE under reducing (Lane 1) and non-reducing (Lane 2) conditions.



The activity of CDK1/Cyclin B1 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.