

**Catalog No.: RP03303LQ** **Recombinant**

Species	Gene ID	Swiss Prot
Human	983&898	P06493&P24864

N-GST (CDK1) & N-His-Flag (Cyclin E1)

CDK1; CDC2; CDC28A; CDKN1; P34CDC2;  
Cyclin-dependent kinase 1; Cyclin E1;  
CCNE1; CCNE; G1/S-specific cyclin-E1

<b>Source</b>	<b>Purification</b>
Baculovirus-Insect Cells	≥ 90 % as determined by SDS-PAGE; ≥ 90 % as determined by HPLC.

## 60.6 kDa/51.0 kDa 50-60 kDa/48-52 kDa

< 1 EU/μg of the protein by LAL method.

Supplied as a 0.22 µm filtered solution in 50 mM Tris-HCl, 150 mM NaCl, 5% glycerol, 1 mM DTT. (pH 7.5). Contact us for customized product form or formulation.

Please use running water to thaw it quickly.

 | 400-999-6126

 | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

CDK1 also known as cyclin-dependent kinase 1 or cell division cycle protein 2 homolog is a highly conserved protein that functions as a serine/threonine protein kinase, and is a key player in cell cycle regulation. CDK1 is a small protein (approximately 34 kDa), and is highly conserved. When bound to its cyclin partners, CDK1 phosphorylation leads to cell cycle progression. Given its essential role in cell cycle progression, CDK1 is highly regulated. Most obviously, CDK1 is regulated by its binding with its cyclin partners. Cyclin binding alters access to the active site of CDK1, allowing for CDK1 activity. Cyclin E1 functions as a regulator of S phase entry and progression in mammalian cells. It was found to associate with, and be involved in, the phosphorylation of NPAT protein (nuclear protein mapped to the ATM locus), which participates in cell-cycle regulated histone gene expression and plays a critical role in promoting cell-cycle progression in the absence of pRB.

Recombinant Human CDK1&Cyclin E1 Protein is produced by Baculovirus-Insect Cells expression system. The target protein is expressed with sequence (Met1-Met297 (CDK1) & Met1-Glu410 (Cyclin E1)) of Human CDK1&CCNE1 (Accession #P06493&P24864) fused with a N-GST (CDK1) & N-His-Flag (Cyclin E1) tag.

The activity of CDK1/Cyclin E1 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.

The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.


Store at -70°C. This product is stable at  $\leq -70^{\circ}\text{C}$  for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature.

Aliquots below 10  $\mu$ L are not advisable. Product must not be stored in diluted solutions. Avoid repeated freeze-thaw cycles.


Avoid repeated freeze/thaw cycles.

For your safety and health, please wear a lab coat and disposable gloves for handling.

Validation Data



Recombinant Human CDK1&Cyclin E1 Kinase was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.



The activity of CDK1/Cyclin E1 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.