

**Catalog No.: RP03309LQ** **Recombinant**

Species	Gene ID	Swiss Prot
Human	1017&890	P24941&P20248

N-GST (CDK2) & N-His (Cyclin A2)

CDK2; CDKN2; Cyclin-dependent kinase 2; Cyclin A2; CCNA2; CCN1; CCNA; Cyclin-A2

Source	Purification
Baculovirus-Insect Cells	≥ 90 % as determined by SDS-PAGE; ≥ 90 % as determined by HPLC.

60.4 kDa/32.8 kDa	50-60 kDa/25-30 kDa
-------------------	---------------------

< 1 EU/μg of the protein by LAL method.

Supplied as a 0.22 µm filtered solution in 20 mM HEPES, 0.20 mM NaCl, 5% glycerol, 1 mM DTT. (pH 7.5). Contact us for customized product form or formulation.

Please use running water to thaw it quickly.

 | 400-999-6126

✉ | [cn.market@abclonal.com.cn](mailto:cn.market@abclonal.com.cn)

 | [www.abclonal.com.cn](http://www.abclonal.com.cn)

CDK2, also known as cyclin-dependent kinase 2, or cell division protein kinase 2, is a member of the cyclin-dependent kinase family of Ser/Thr protein kinases. This protein associates with and is regulated by the regulatory subunits of the complex including cyclin E or A. Cyclin E binds G1 phase CDK2, which is required for the transition from G1 to S phase while binding with Cyclin A is required to progress through the S phase. Abnormal expression of cyclin A2 is associated with chromosomal instability and tumor proliferation, while inhibition leads to decreased tumor growth. Therefore, CDK2 and its cyclin binding partners represent possible therapeutic targets for new cancer therapeutics.

Recombinant Human CDK2&Cyclin A2 Protein is produced by Baculovirus-Insect Cells expression system. The target protein is expressed with sequence (Met1-Leu298 (CDK2) & Glu174-Leu432 (Cyclin A2)) of Human CDK2&CCNA2 (Accession #P24941&P20248) fused with a N-GST (CDK2) & N-His (Cyclin A2) tag.


The activity of CDK2/Cyclin A2 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.

Store at -70°C. This product is stable at  $\leq -70^{\circ}\text{C}$  for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature.


Aliquots below 10  $\mu$ L are not advisable. Product must not be stored in diluted solutions. Avoid repeated freeze-thaw cycles.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human CDK2&Cyclin A2 Kinase was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.



The activity of CDK2/Cyclin A2 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.