

# Recombinant Human CDK2&Cyclin E1 Kinase

Catalog No.: RP03310LQ Recombinant

## **Sequence Information**

 Species
 Gene ID
 Swiss Prot

 Human
 1017&898
 P24941&P24

 864
 864

#### **Tags**

N-GST (CDK2) & N-His-Flag (Cyclin E1)

#### **Synonyms**

CDK2; CDKN2; Cyclin-dependent kinase 2; Cyclin E1; CCNE1; CCNE; G1/S-specific cyclin-E1

## **Product Information**

Source	Purification
Baculovirus-Insect	≥ 90 % as
Cells	determined by SDS-
	PAGE;≥ 90 % as
	determined by
	HPLC.

## Calculated MW Observed MW

60.5 kDa/51.1 kDa 50-60 kDa/48-52

kDa

#### **Endotoxin**

< 1 EU/µg of the protein by LAL method.

#### **Formulation**

Supplied as a 0.22  $\mu$ m filtered solution in 50 mM Tris-HCl, 200 mM NaCl, 5% glycerol. (pH 7.5). Contact us for customized product form or formulation.

#### Reconstitution

Please use running water to thaw it quickly.

#### Contact

<u>a</u>		400-999-6126
$\bowtie$		cn.market@abclonal.com.cn
•	Т	www.abclonal.com.cn

## **Background**

CDK2, also known as cyclin-dependent kinase 2, or cell division protein kinase 2, is a member of the cyclin-dependent kinase family of Ser/Thr protein kinases. This protein associates with and is regulated by the regulatory subunits of the complex including cyclin E or A. Cyclin E binds G1 phase CDK2, which is required for the transition from G1 to S phase while binding with Cyclin A is required to progress through the S phase. The cyclin E protein contains a section called the cyclin box, which interacts with the PSTAIRE helix on CDK2 to enact a conformational change in CDK2's T loop. Once the CDK2/Cyclin E1 dimer is formed and activated, it phosphorylates several important proteins including proteins involved in centrosome duplication (NPM, CP110, Mps1), DNA synthesis (Cdt1), DNA repair (Brca1, Ku70), histone gene transcription (p220/NPAT, CBP/p300, HIRA) and Cdk inhibitors.

## **Basic Information**

#### Description

Recombinant Human CDK2&Cyclin E1 Protein is produced by Baculovirus-Insect Cells expression system. The target protein is expressed with sequence (Met1-Leu298 (CDK2) & Met1-Glu410 (Cyclin E1)) of Human CDK2&CCNE1 (Accession #P24941&P24864) fused with a N-GST (CDK2) & N-His-Flag (Cyclin E1) tag.

#### **Bio-Activity**


The activity of CDK2/Cyclin E1 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.

#### **Storage**


Store at  $-70^{\circ}$ C. This product is stable at  $\leq -70^{\circ}$ C for up to 1 year from the date of receipt. For optimal storage, aliquot into smaller quantities after centrifugation and store at recommended temperature.

Aliquots below 10  $\mu\text{L}$  are not advisable. Product must not be stored in diluted solutions. Avoid repeated freeze-thaw cycles.

Avoid repeated freeze/thaw cycles.



Recombinant Human CDK2&Cyclin E1 Kinase was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.



The activity of CDK2/Cyclin E1 is based on the MSA technology, and the content and ratio of the substrate and the product are directly separated and detected in real time and dynamically by the different migration rates of the substrate and the product after the enzymatic reaction.