

Human Glycoprotein IIb/IIIa Receptor Monoclonal Antibody

Catalog No.: YR0105

Basic Information

Molecular Weight

150 kDa

Endotoxin

<1EU/mg (<0.001EU/ μ g)Determined by LAL gel clotting assay

Sterility

0.2 μm filtration

Aggregation

<5% Determined by SECP

Purity

>95% Determined by SDS-PAGE

Background

Abciximab Biosimilar uses the same protein sequences as the therapeutic antibody abciximab. Abciximab, a glycoprotein IIb/IIIa receptor antagonist, is a platelet aggregation inhibitor mainly used during and after coronary artery procedures like angioplasty to prevent platelets from sticking together and causing thrombus (blood clot) formation within the coronary artery. It is a glycoprotein IIb/IIIa inhibitor. While abciximab has a short plasma half-life, due to its strong affinity for its receptor on the platelets, it may occupy some receptors for weeks. In practice, platelet aggregation gradually returns to normal about 96 to 120 hours after discontinuation of the drug. Abciximab is made from the Fab fragments of an immunoglobulin that targets the glycoprotein IIb/IIIa receptor on the platelet membrane.

Reported Applications

ELISA, neutralization, functional assays such as bioanalytical PK and ADA assays, and those assays for studying biological pathways

Immunogen Information

Clone

Isotype Human IgG1 kappa

Immunogen

glycoprotein (GP) IIb/IIIa

Abciximab Biosimilar

RecommendedIsotype Control(s)

In Vivo Grade Recombinant Human IgG1 Kappa Isotype Control Antibody

Recommended Dilution Buffer

1×PBS pH 7.4

Contact

<u>a</u>		400-999-6126
\bowtie	Τ	cn.market@abclonal.com.cn
•	ī	www.abclonal.com.cn

Product Information

Production

Purification

Purified from cell culture supernatant in an animal-free facility

Protein A or G purification

Storage

Store at 2 - 8°C. 2 - 8°C for up to 4 weeks and -80°C for long term storage (Avoid repeated freezing and thawing)