

Human CD4 Monoclonal Antibody

Catalog No.: YR0108

Basic Information

Molecular Weight

150 kDa

Endotoxin

<1EU/mg (<0.001EU/μg) Determined by
LAL gel clotting assay

Sterility

0.2 μm filtration

Aggregation

<5% Determined by SECP

Purity

>95% Determined by SDS-PAGE

Background

CD4 is present on a subset of T lymphocytes ("helper/inducer" T cells) and is also expressed at a lower level on monocytes, tissue macrophages and granulocytes. The antigen is involved in binding to MHC class II molecules. The intracellular domain of the antigen is associated with p56lck protein tyrosine kinase.

Reported Applications

ELISA, neutralization, functional assays such as bioanalytical PK and ADA assays, and those assays for studying biological pathways

Immunogen Information

Clone

Clenoliximab Biosimilar

Isotype

Human IgG4 kappa

Immunogen

Human CD4

Recommended Isotype Control(s)

In Vivo Grade Recombinant Human IgG4-S228P Kappa Isotype Control Antibody


Recommended Dilution Buffer

1×PBS pH 7.4

Contact

 | 400-999-6126

 | cn.market@abclonal.com.cn

 | www.abclonal.com.cn

Product Information

Production

Purified from cell culture supernatant in an animal-free facility

Purification

Protein A or G purification

Storage

Store at 2 - 8°C. 2 - 8°C for up to 4 weeks and -80°C for long term storage (Avoid repeated freezing and thawing)