Human CCR4 Monoclonal Antibody

Catalog No.: YR0126



Basic Information

Molecular Weight

150 kDa

Endotoxin

<1EU/mg (<0.001EU/ μ g)Determined by LAL gel clotting assay

Sterility

0.2 µm filtration

Aggregation

<5% Determined by SECP

Purity

>95% Determined by SDS-PAGE

Background

Mogamulizumab is a humanized monoclonal antibody (mAb) directed against CC chemokine receptor 4 (CCR4) for the treatment of Mycosis Fungoides (MF) and Sézary Syndrome (SS), the most common subtypes of cutaneous T-cell lymphoma. Cutaneous T-cell lymphomas occur when certain white blood cells, called T cells, become cancerous; these cancers typically affect the skin, causing various types of skin lesions. Mogamulizumab selectively binds to and inhibits the activity of CCR4, which may block CCR4-mediated signal transduction pathways and, so, chemokine-mediated cellular migration and proliferation of T cells, as well as chemokine-mediated angiogenesis. Additionally, this agent may induce antibody-dependent cell-mediated cytotoxicity (ADCC) against CCR4-positive T cells. CCR4, a G-coupled-protein receptor for C-C chemokines such MIP-1, RANTES, TARC and MCP-1, is expressed on the surfaces of some types of T cells, endothelial cells, and certain types of neurons. CCR4, also known as CD194, may be overexpressed on adult T-cell lymphoma (ATL) and peripheral T-cell lymphoma (PTCL) cells. In addition to directly targeting malignant T cells expressing CCR4, mogamulizumab depletes Treg cells, an important therapeutic target in many human cancers because of their role in suppressing host antitumor immunity.

Reported Applications

ELISA,neutralization,functional assays such as bioanalytical P K and ADA assays,and those assays for studying biological pathways

Immunogen Information

Clone

Mogamulizumab Biosimilar

Isotype

Human IgG1 kappa

Immunogen

Human CCR4

RecommendedIsotype Control(s)

In Vivo Grade Recombinant Human IgG1 Kappa Isotype Control Antibody

Recommended Dilution Buffer

1×PBS pH 7.4

Contact

| 6 | 400-999-6126 |
|-----------|---------------------------|
| \bowtie | cn.market@abclonal.com.cn |
| • | www.abclonal.com.cn |

Product Information

Production

Purification

Purified from cell culture supernatant in an animal-free facility

Protein A or G purification

Storage

Store at 2 - 8°C. 2 - 8°C for up to 4 weeks and -80°C for long term storage (Avoid repeated freezing and thawing)