

Liraglutide Biosimilar

Catalog No.: YR0141

Basic Information

Molecular Weight

3.8 kDa

Endotoxin

<1EU/mg (<0.001EU/μg) Determined by LAL gel clotting assay

Sterility

0.2 μm filtration

Aggregation

<5% Determined by SECP

Purity

>95% Determined by SDS-PAGE

Background

Liraglutide, a recombinant peptide which presents 97% homology with human glucagon-like peptide-1 (GLP-1), differing primarily by substituting arginine for lysine at position 34. Liraglutide is made by attaching a C-16 fatty acid (palmitic acid) with a glutamic acid spacer on the remaining lysine residue at position 26 of the peptide precursor. Liraglutide is resistant to the dipeptidylpeptidase-4 enzyme which degrades the natural hormone. As a glucagon-like peptide-1 receptor agonist (GLP-1 receptor agonist) also known as incretin mimetics, Liraglutide activates the GLP-1 receptor and exerts an incretin mimetic effect during at least 24 hours after a single subcutaneous injection. Besides a glucose-dependent stimulatory effect of insulin secretion, liraglutide inhibits glucagon secretion and retards gastric emptying. Liraglutide is an anti-diabetic medication used to treat type 2 diabetes, obesity, and chronic weight management. Liraglutide is given by injection under the skin. Liraglutide works by increasing insulin release from the pancreas and decreases excessive glucagon release.

Reported Applications

ELISA, neutralization, functional assays such as bioanalytical PK and ADA assays, and those assays for studying biological pathways

Immunogen Information

Clone

Liraglutide

Isotype

Immunogen

Recommended Isotype Control(s)

Recommended Dilution Buffer

1×PBS pH 7.0

Contact

☎ | 400-999-6126

✉ | cn.market@abclonal.com.cn

🌐 | www.abclonal.com.cn

Product Information

Production

Purified from cell culture supernatant in an animal-free facility

Purification

Protein A or G purification

Storage

2 - 8°C for up to 4 weeks and -80°C for long term storage (Avoid repeated freezing and thawing)