

In Vivo Grade Recombinant Human IgG2 Isotype Control Antibody

Catalog No.: YR0178

Basic Information

Molecular Weight

150 kDa

Endotoxin

<1EU/mg (<0.001EU/ μ g)Determined by LAL gel clotting assay

Sterility

0.2 µm filtration

Aggregation

<5% Determined by SECP

Purity

>95% Determined by SDS-PAGE

Background

Isotype control antibodies (negative control antibodies or reference antibodies) are frequently used in various antibody-related in vivo and in vitro experiments such as ELISA, Western blot, immunofluorescence, immunohistochemistry, immunoprecipitation, and flow cytometry. Isotype control antibodies are used as negative controls of primary antibodies to check presence of non-specific binding to Fc receptors on the cell surface, other cellular proteins, carbohydrates, and lipids. Thus, the isotype control antibody normally matches the host species and isotype, including light chains (such as human IgG1 kappa antibody), concentration, and conjugation format of the experimental primary antibody.

Reported Applications

an isotype-

matched negative control use d in ELISA, Western Blot (WB), Flow Cytometry (Flow), Immuno precipitation (IP), Immunohistoc hemistry (Paraffin) (IHC (P)), I mmunohistochemistry (Frozen) (IHC (F)), and in vivo animal model research

Immunogen Information

Clone 4F17 Isotype

Human IgG2 kappa

Immunogen

RecommendedIsotype Control(s)

Recommended Dilution Buffer

1×PBS

Contact

<u>a</u>		400-999-6126
\bowtie	Τ	cn.market@abclonal.com.cn
•	ī	www.abclonal.com.cn

Product Information

Production

Purification

Purified from cell culture supernatant in an animal-free facility

Protein A or G purification

Storage

Store at $2 - 8^{\circ}$ C. $2 - 8^{\circ}$ C for up to 4 weeks and -80° C for long term storage (Avoid repeated freezing and thawing)