Leader in Biomolecular Solutions for Life Science

www.abclonal.com

Rabbit anti-Human IgG Specific Monoclonal Antibody, clone 1F1 (minimum binding to Monkey/Mouse/Rat IgG)

Catalog No.: YR0505

Basic Information

Molecular Weight
150kDa

Endotoxin
<1EU/mg (<0.001EU/μg)Determined by LAL gel clotting assay

Sterility
0.2 μm filtration

Aggregation
<5% Determined by SECP

Purity
>95% Determined by SDS-PAGE

Background

Rabbit anti-Human IgG Specific Monoclonal Antibody clone 1F1 is a recombinant rabbit Monoclonal antibody, that recognizes human immunoglobulin isotypes (IgG1, IgG2, IgG3, IgG4), total IgG. It exhibits minimum binding Cynomolgus macaque total IgG, Rhesus macaque total IgG, Marmoset total IgG, Mouse total IgG, Rat total IgG.

Reported Applications

ELISA,SPR

Immunogen Information

Clone
1F1

Isotype
Rabbit IgG1

Immunogen

RecommendedIsotype Control(s)

Recommended Dilution Buffer
1×PBS pH 7.3

Contact

400-999-6126

cn.market@abclonal.com.cn

www.abclonal.com.cn


Product Information

Production
Purified from cell culture supernatant in an animal-free facility


Purification
Protein A/G

Storage
Store at -20℃


Validation Data




Regular ELISA binding curve demonstrating the recognition of Rabbit anti-Human IgG Specific Monoclonal Antibody. A microtiter plate was coated overnight with various antibodies at a concentration of 2 μg/mL. After washing and blocking with PBST + 5% skim milk, increasing concentrations of the Rabbit Anti-Human IgG Fab Antibody (clone 1F1) was added. Horseradish Peroxidase conjugated goat anti-rabbit IgG (Jackson, Cat.: 111-035-045) was used for final detection.




Regular ELISA binding curve demonstrating the recognition of Rabbit anti-Human IgG Specific Monoclonal Antibody. A microtiter plate was coated overnight with various antibodies at a concentration of 2 μg/mL. After washing and blocking with PBST + 5% skim milk, increasing concentrations of the Rabbit Anti-Human IgG Fab Antibody (clone 1F1) was added. Horseradish Peroxidase conjugated goat anti-rabbit IgG (Jackson, Cat.: 111-035-045) was used for final detection.




Capture ELISA binding curve demonstrating the recognition of Rabbit anti-Human IgG Specific Monoclonal Antibody. A microtiter plate was coated overnight with Rabbit anti-Human IgG Specific Monoclonal Antibody (clone 1F1) at a concentration of 2 μg/mL. After washing and blocking with PBST + 5% skim milk, increasing concentrations of various biotinylated antibodies was added. Horseradish Peroxidase conjugated Neutravidin (Thermo Fisher, 31001) was used for final detection.




Capture ELISA binding curve demonstrating the recognition of Rabbit anti-Human IgG Specific Monoclonal Antibody. A microtiter plate was coated overnight with Rabbit anti-Human IgG Specific Monoclonal Antibody (clone 1F1) at a concentration of 2 μg/mL. After washing and blocking with PBST + 5% skim milk, increasing concentrations of various biotinylated antibodies was added. Horseradish Peroxidase conjugated Neutravidin (Thermo Fisher, 31001) was used for final detection.




A microtiter plate was coated overnight with VEGF at a concentration of 2 μg/mL. After washing and blocking with PBST + 5% skim milk, 10% cynomolgus macaque serum was added spiked with increasing concentrations of bevacizumab (hIgG1 kappa)(from 10 μg/mL). Clone 1F1 was added. Another unrelated rabbit mAb used for negative control. Detection was performed using HRP conjugated goat anti-rabbit 2nd Ab. No cynomolgus serum added sample is used as control.



Rabbit anti-Human IgG Specific Monoclonal Antibody (Clone 1F1) captured on C5 Chip can bind Human IgG1 with an affinity constant of 10.42 nM as determined in SPR assay.





Validation Data

Rabbit anti-Human IgG Specific Monoclonal Antibody (Clone 1F1) captured on C5 Chip can bind Human IgG3 with an affinity constant of 18.26 nM as determined in SPR assay

Rabbit anti-Human IgG Specific Monoclonal Antibody (Clone 1F1) captured on C5 Chip can bind Human IgG4 with an affinity constant of 5.07 nM as determined in SPR assay.